Crices

prior to administration of the liposome composition and wherein the liposome has an average diameter of from at least about 200 nm to about 5000 nm.

 $\frac{2}{\sqrt{2}}$ 

- 14. (Amended) The method of Claim 13, wherein the functional group is a hydroxyl, thiol epoxide or amine group.
- 16. (Amended) The method of Claim 15, wherein the bioactive agent is a contrast agent, antibacterial agent, antiviral agent, antifungal agent, anti-parasitic agent, tumoricidal agent, carbohydrate, polypeptide, peptide, toxin, enzyme, hormone, neurotransmitter, glycoprotein, lipoprotein, immunomodulator, vasodilator, dye, radiolabel, fluorescent compound, receptor binding molecule, anti-inflammatory agent, mydriatic compound, local anesthetic, narcotic, vitamin, polynucleotide, nucleoside, or nucleotide.

Please cancel Claims 18-53, without prejudice to or disclaimer of the subject matter contained therein.

Please add new Claims 54-57:

(New) The method of Claim 16, wherein the contrast agent is a MRI contrast agent, a radio-contrast agent, or a water-soluble iodinated contrast agent.

04

- 55. (New) The method of Claim 15, wherein the bioactive agent is an antimetabolite.
- 2019 56. (New) The method of Claim 15, wherein the bioactive agent is an immunoglobulin.

Application No. 09/398,934 Attorney's Docket No. 033364-280 Page 3

Corold.

2 57. (New) The method of Claim 15, wherein the bioactive agent is a radio-opaque compound.